

August 1991

From Virasoro Constraints in Kontsevich's Model to $\mathcal{W}$ -constraints in 2-matrix Models

A.Marshakov, A.Mironov

Department of Theoretical Physics

P.N.Lebedev Physical Institute

*Leninsky prospect, 53, Moscow, 117 924*¹,

A.Morozov

Institute of Theoretical and Experimental Physics,

Bol.Cheremushkinskaya st., 25, Moscow, 117 259

ABSTRACT

The Ward identities in Kontsevich-like 1-matrix models are used to prove at the level of discrete matrix models the suggestion of Gava and Narain, which relates the degree of potential in asymmetric 2-matrix model to the form of $\mathcal{W}$ -constraints imposed on its partition function.

¹E-mail address: theordep@sci.fian.msk.su

1 Introduction

Matrix models, originally developed [1] as an alternative approach to (so far 2-dimensional) quantum gravity, are now understood to possess a deep and interesting mathematical structure of their own. Indeed, their partition functions are usually subjected to infinite sets of constraints, which can be formulated in a form of differential equations with respect to the "time"-variables, and as a corollary they appear proportional to " τ -functions" of integrable hierarchies [2].

While this general scheme is more or less well understood at the level of discrete 1-matrix models [3,4], it is still in many respects an open question when continuum limit and/or *multimatrix* models are concerned. An important breakthrough in the study of continuum limit of 1-matrix models is due to a recent conjecture of M.Kontsevich [5], who in fact suggested that the proper continuum limit of the Hermitean 1-matrix model can be described in terms of a somewhat different matrix model, to be referred later as Kontsevich's model. Its partition function is essentially proportional to the $N \times N$ (anti)Hermitean matrix integral

$$\mathcal{F}\{\Lambda\} \equiv \int DX \exp(-trW[X] + tr\Lambda X) \quad (1)$$

with $W[X] = X^3$ and $N \rightarrow \infty$. This conjecture was strongly supported by a recent proof [6] that partition function of Kontsevich's model satisfies exactly the same set of differential equations (Virasoro constraints [2]) as continuum limit of the original Hermitean 1-matrix model (so that it remains only to understand what are the requirements, which guarantee the uniqueness of the solution of these Virasoro constraints). In [1] it was shown that Virasoro constraints arise from the obvious Ward identities, satisfied by $\mathcal{F}(\Lambda)$,

$$(tr\Lambda^p W'[\partial/\partial\Lambda_{tr}] - tr\Lambda^{p+1})\mathcal{F}\{\Lambda\} = 0 \quad (2)$$

as a result of a change (Miwa transformation) of the argument

$$\Lambda \rightarrow \{T_m = \frac{1}{m + \frac{1}{2}} tr\Lambda^{-m-\frac{1}{2}} - \frac{4}{3\sqrt{3}}\delta_{m,1}\} . \quad (3)$$

In its turn, eq.(2) states that the integral (1) is invariant under infinitesimal shift of integration variable,

$$X \rightarrow X + \Lambda^p . \quad (4)$$

The implication of Kontsevich's conjecture in this context (for more physical implication see ref.[6]) is that the Miwa transformation (3) allows us to substitute a sophisticated double-scaling limit in conventional matrix models [1] by a naive limit of $N \rightarrow \infty$ in the model like (1). This opens promising possibilities in the study of continuum matrix models.

Somewhat unexpectedly this fresh view on continuum limits provides also a new approach to the study of *discrete* multimatrix models. One of the main problems about them is the lack of understanding (and even a derivation) of differential equations which substitute Virasoro constraints of 1-matrix models

(and are believed [2] to be expressible in terms of generators of $\mathcal{W}$ -algebras). What we are going to demonstrate in this letter is that just the same Ward identities (2) for the matrix integral (1) provide a simple proof of the $\mathcal{W}$ -like constraints in a discrete (i.e. at finite N) 2-matrix model.

Moreover, the recent observation due to E.Gava and K.Narain [7], stating that the spin of $\mathcal{W}$ -constraint in fact coincides with the power of potential $W[X]$ is naturally explained in this way. (This suggestive result has been obtained in [7] by a tedious examination of continuum limit of loop equations for specific 2-matrix model with $W[X] = X^3$).

2 The main results

The partition function of the discrete Hermitean 2-matrix model [8] is given by a double integral over $N \times N$ Hermitean matrices X and Λ :

$$Z_{V,W} = \int DX D\Lambda \exp(-tr\{V[\Lambda] + W[X] - \Lambda X\}) \equiv \int D\Lambda e^{-trV[\Lambda]} \mathcal{F}_W\{\Lambda\} . \quad (5)$$

The potentials V and W are conventionally parameterized by the corresponding time-variables

$$\begin{aligned} V[\Lambda] &= \sum_{k \geq 0} t_k \Lambda^k , \\ W[X] &= \sum_{k \geq 0} s_k X^k \end{aligned} \quad (6)$$

and the partition function $Z_{V,W}$ is usually treated as a functional of $\{t_k\}$ and $\{s_k\}$. Below we will use the obvious notation $Z_{V,W} \equiv Z_W\{t_k\}$. While in [2] it was suggested that the continuum (*i.e.* $N \rightarrow \infty$) limit of $Z_{V,W}$ in the case of $W = V$ (this $K = \infty$) is annihilated by a set of operators which form $\mathcal{W}_3$ -algebra, the results of ref.[7] imply, at least, that the structure of constraints in the asymmetric situation $W \neq V$ is more complicated: generators of $\mathcal{W}_K$ -algebra (expressed in terms of t -variables) annihilate $Z_W\{t_k\}$ whenever $W[X]$ is a polynomial of power K . Our purpose below is to explore the origin of this important phenomenon in the most transparent way.

The natural derivation arises from comparison of eqs. (5) and (2). Indeed:

$$\frac{\partial Z_W}{\partial t_{p+1}} = - \int D\Lambda e^{-trV[\Lambda]} tr \Lambda^{p+1} \mathcal{F}\{\Lambda\} = - \int D\Lambda e^{-trV[\Lambda]} tr(\Lambda^p W'[\frac{\partial}{\partial \Lambda_{tr}}]) \mathcal{F}\{\Lambda\} . \quad (7)$$

After integration by parts it turns into

$$\int D\Lambda \mathcal{F}\{\Lambda\} tr(W'[-\frac{\partial}{\partial \Lambda_{tr}}] \Lambda^p) e^{-trV[\Lambda]} = \sum_{k > 0}^K k s_k \int D\Lambda \mathcal{F}\{\Lambda\} tr((-\frac{\partial}{\partial \Lambda_{tr}})^{k-1} \Lambda^p) e^{-trV[\Lambda]} . \quad (8)$$

The leading term in the sum on the r.h.s., i.e.

$$\begin{aligned}
& K s_K \int D\Lambda \mathcal{F}\{\Lambda\} \{tr \Lambda^p (V'[\Lambda])^{K-1} + O(V^{K-2})\} = \\
& = K s_K \sum_{a_1, \dots, a_{K-1}} a_1 t_{a_1} \dots a_{K-1} t_{a_{K-1}} \frac{\partial}{\partial t_{a_1 + \dots + a_{K-1} + p + 1 - K}} Z_W\{t\}
\end{aligned} \tag{9}$$

is a “classical”² piece of the operator $\mathcal{W}_{p+1-K}^{(K)}$ — $(p + K - 1)$ -th harmonic of the spin- K generator of $\mathcal{W}_K$ -algebra — acting on $Z_W\{t\}$. (Note also that according to (8) $p \geq 0$). This explains essentially the very origin of $\mathcal{W}_K$ -constraint and its intimate relation to the form of potential $W[\Lambda]$ in complete agreement with [7].

In general, the Ward identity (7) can be rewritten as a set of constraints

$$\left(-\frac{\partial}{\partial t_{p+1}} + \sum_{k \geq 0}^K k s_k \tilde{\mathcal{W}}_{p+1-k}^{(k)}\{t\} \right) Z_W\{t\} = 0 . \tag{10}$$

Operators $\tilde{\mathcal{W}}^{(k)}$ are defined by

$$\tilde{\mathcal{W}}_{p+1-k}^{(k)} e^{-tr V[\Lambda]} = -tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} \Lambda^p \right) e^{-tr V[\Lambda]} . \tag{11}$$

They obey recurrent relation

$$\tilde{\mathcal{W}}_p^{(k+1)} = \sum_n n t_n \tilde{\mathcal{W}}_{n+p}^{(k)} + \sum_{a+b=p+k-1} \frac{\partial}{\partial t_a} \tilde{\mathcal{W}}_{b+1-k}^{(k)} ; p \geq -k \tag{12}$$

with

$$\tilde{\mathcal{W}}_p^{(2)} = \mathcal{L}_p = \sum_n n t_n \frac{\partial}{\partial t_{n+p}} + \sum_{a+b=p} \frac{\partial^2}{\partial t_a \partial t_b} ; p \geq -1 , \tag{13}$$

$$\tilde{\mathcal{W}}_p^{(1)} = \mathcal{J}_p = \frac{\partial}{\partial t_p} ; p \geq 0 . \tag{14}$$

In order to derive (12) one has to take explicitly one derivative $\partial/\partial \Lambda_{tr}$ in (11):

$$\begin{aligned}
& \tilde{\mathcal{W}}_p^{(k+1)} e^{-tr V[\Lambda]} = \\
& = tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^k \Lambda^{p+k} \right) e^{-tr V[\Lambda]} = tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} \left(-\frac{\partial}{\partial \Lambda_{tr}} \right) \Lambda^{p+k} \right) e^{-tr V[\Lambda]} = \\
& = tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} V'(\Lambda) \Lambda^{p+k} \right) e^{-tr V[\Lambda]} - \sum_{a+b=p+k-1} tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} \Lambda^b \right) tr \Lambda^a e^{-tr V[\Lambda]} = \\
& = \sum_n n t_n tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} \Lambda^{p+n+k-1} \right) e^{-tr V[\Lambda]} + \\
& + \sum_{a+b=p+k-1} \frac{\partial}{\partial t_a} \left(tr \left(\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{k-1} \Lambda^b \right) e^{-tr V[\Lambda]} \right) = \\
& = \left(\sum_n n t_n \tilde{\mathcal{W}}_{n+p}^{(k)} + \sum_{a+b=p+k-1} \frac{\partial}{\partial t_a} \tilde{\mathcal{W}}_{b+1-k}^{(k)} \right) e^{-tr V[\Lambda]} .
\end{aligned} \tag{15}$$

²In the sense of ref.[9]

Eqs.(13) and, especially, (14) are trivially derived. Eqs.(12) and (13) imply that all the $\tilde{\mathcal{W}}^{(k)}$ -operators are in fact proportional to linear combinations of Virasoro operators $\mathcal{L} = \tilde{\mathcal{W}}^{(2)}$. This may explain how in the continuum limit a single Ward identity (10) can give rise to entire set of constraints with lower spins. However, this topic is beyond the scope of this Letter. The continuum limit of these eqs. may be studied along the lines of [7].

3 Formulation in terms of [9,10]

While from the point of view of derivations it is illuminating first to study the simplest Ward identity (3) in the “1-matrix component” of 2-matrix model, and then apply it to the (more sophisticated) analysis of 2-matrix case, it is of course possible to treat the resulting $\mathcal{W}$ -constraints (10) as Ward identities in the entire 2-matrix model, related to the following infinitesimal change of integration variables

$$\begin{aligned}\delta X &= \Lambda^p, p \geq 0 \\ \delta \Lambda &= \left(\sum_{m=0}^K m s_m \sum_{k=0}^{m-2} (-)^{k+1} (V')^k X^{m-2-k} \right) \Lambda^p + \text{”quantum corrections”} .\end{aligned}\tag{16}$$

This variation of variables induces the variation of potential:

$$\delta S = \left(\sum_{m=0}^K m s_m (-)^m (V')^{m-1} \right) \Lambda^p + \Lambda^{p+1} + \text{”quantum corrections”} .\tag{17}$$

The first term in this expression gives rise to the ”classical” part of $\tilde{\mathcal{W}}$ -algebra and the second one produces the derivative $\partial/\partial t_{p+1}$ in (10).

While the X -component of the variation (16) (which is of course nothing but (3)) does not change the integration measure $DXD\Lambda$, this is not true for Λ -component. The corresponding Jacobian is responsible for the “quantum” contributions to (16) and (17).

Eqs. (16)-(17) provide a possible generalization to the 2-matrix case of the derivation [9,10] of the Virasoro constraints in the discrete 1-matrix model from the Ward identities associated with the shift $M \rightarrow M + \epsilon M^{n+1}$ ($n \geq -1$) of the integration variable.

4 Some results about $\tilde{\mathcal{W}}$ -operators

4.1 A short summary

The purpose of this section is to begin the systematic study of $\tilde{\mathcal{W}}$ -constraints.

First of all we should stress that eq.(11) defines only *positive* ($p \geq 1 - K$, to be exact) harmonics $\tilde{\mathcal{W}}_p^{(K)}$ of $\tilde{\mathcal{W}}^{(K)}$ -operators. We do not address the structure of entire $\tilde{\mathcal{W}}^{(K)}$ -operators below.

Second, an important question is whether the set of $\tilde{\mathcal{W}}$ -constraints (10) is closed. We shall demonstrate in sect.4.3 below that this is indeed the case. Namely it will be proved that

$$[\tilde{\mathcal{W}}_p^{(K)}, \tilde{\mathcal{W}}_q^{(K)}] \in \text{Span} \tilde{\mathcal{W}}_r^{(K)}, \quad (18)$$

where the notation *Span* implies all possible linear combinations with coefficients linearly depending of times and derivatives and $r \geq 1 - K$ as long as $p, q \geq 1 - K$. (In order to complete the proof one should also demonstrate that the commutators of $\tilde{\mathcal{W}}_p^{(K)}$ with lower-spin operators $\tilde{\mathcal{W}}^{(n)}$, $n < K$ are appropriately adjusted. This piece of the proof is omitted from sect.4 at the moment: we do not know any elegant way to represent the entire algebra of $\tilde{\mathcal{W}}$ -operators.) It deserves noting that (18) remains a highly non-trivial property of $\tilde{\mathcal{W}}$ -algebra: by no means it follows from the fact that $\tilde{\mathcal{W}}_p^{(K)}$ may be expressed through $\mathcal{L}_q$ or $\mathcal{J}_q$. Moreover $\tilde{\mathcal{W}}^{(K)}Z = 0$ does not imply that lower $\tilde{\mathcal{W}}^{(n)}$, $n < K$ annihilate Z , therefore (18) is absolutely crucial for the closeness of the set of constraints (11). The fact that this set is indeed closed makes an interesting exercise to observe the appearance of the entire tower of $\tilde{\mathcal{W}}^{(n)}$ -constraints (with all spins $n \leq K$) in the continuum limit from the single spin- K constraint at the discrete level.

Third, just the same $\tilde{\mathcal{W}}$ -operators were found in [6] in a somewhat different context. In sect 4.5 we shall prove that they are really the same, thus demonstrating a kind of universal nature of $\tilde{\mathcal{W}}$ -operators, at least, in the framework of discrete matrix models.

Fourth, the fact that the commutator of $\tilde{\mathcal{W}}^{(K)}$ -operators in (18) is not just proportional to $\tilde{\mathcal{W}}_{p+q}^{(2K-2)}$ demonstrates that $\oplus_K \tilde{\mathcal{W}}^{(K)}$ is *not* a Lie algebra (to make it similar to $\mathcal{W}_\infty$, at least, the basis should be changed). This makes $\tilde{\mathcal{W}}$ even more similar to conventional $\mathcal{W}$ -algebras which are also non-linear and closed as soon as only operators $\tilde{\mathcal{W}}^{(n)}$ of spins $n \leq K$ are considered.

Thus the subject of $\tilde{\mathcal{W}}$ -algebras may deserve further investigation.

4.2 Other formulations of $\tilde{\mathcal{W}}$ -operators

Before we turn to the derivation of (18) let us present an explicit expressions for the first members of the $\tilde{\mathcal{W}}$ -family

$$\tilde{\mathcal{W}}^{(2)} = \tilde{\mathcal{L}}_n = \sum_{k>0} kt_k \frac{\partial}{\partial t_{k+n}} + \sum_{a+b=n} \frac{\partial^2}{\partial t_a \partial t_b} \quad (19)$$

$$\begin{aligned} \tilde{\mathcal{W}}_{p-2}^{(3)} = & \sum_{k,l>0} kt_k lt_l \frac{\partial}{\partial t_{p+k+l-2}} + \sum_{k>0} kt_k \sum_{a+b=k+p-2} \frac{\partial^2}{\partial t_a \partial t_b} + \\ & + \sum_{k>0} kt_k \sum_{a+b=p-1} \frac{\partial^2}{\partial t_a \partial t_{k+b-1}} + \sum_{a+b+c=p-2} \frac{\partial^3}{\partial t_a \partial t_b \partial t_c} + \frac{p(p-1)}{2} \frac{\partial}{\partial t_p} \end{aligned} \quad (20)$$

For comparison we write down conventional $\mathcal{W}$ -operators:

$$\mathcal{W}_n^{(2)} = \mathcal{L}_n = \sum_{k>0} kt_k \frac{\partial}{\partial t_{k+n}} + \frac{1}{2} \sum_{a+b=n} \frac{\partial^2}{\partial t_a \partial t_b} \quad (21)$$

$$\mathcal{W}_p^{(3)} = 3 \sum_{k,l>0} kt_k lt_l \frac{\partial}{\partial t_{p+k+l}} + 3 \sum_{k>0} kt_k \sum_{a+b=k+p} \frac{\partial^2}{\partial t_a \partial t_b} + \sum_{a+b+c=p} \frac{\partial^3}{\partial t_a \partial t_b \partial t_c} \quad (22)$$

These operators can be expressed in terms of free field. Let us introduce also the corresponding current (its negative modes are described by eq.(14)):

$$\mathcal{J}(z) = \sqrt{2}\partial\phi(z) = \sum_{k>0} k t_k z^{k-1} + \sum_{k\geq 0} \frac{1}{z^{k+1}} \frac{\partial}{\partial t_k} = \sum_{k>0} \mathcal{J}_{-k} z^{k-1} + \sum_{k\geq 0} \mathcal{J}_k z^{-k-1} \quad (23)$$

Now $\mathcal{W}$ -operators can be described as

$$\mathcal{W}_p^K =: \mathcal{J}^K :_{-K-3} \quad (24)$$

where standard normal ordering $: \dots :$ implies that all positive current modes should be pulled to the left and commutators should be thrown out.

Alternatively, standard normal ordering one can introduce another one (we shall denote it as $\ddagger \dots \ddagger$) which permits one to reproduce $\tilde{\mathcal{W}}$ -operators from the same formulas (24). Namely, new ordering implies that the terms with “wrong” order of current modes should be just thrown out. Let us consider two simplest examples. The first is the Virasoro case, where positive generators can be written as a sum of three terms:

$$\mathcal{J}_+ \mathcal{J}_- + \mathcal{J}_- \mathcal{J}_+ + \mathcal{J}_- \mathcal{J}_-. \quad (25)$$

Then the standard normal ordering leads to the expression (21):

$$\mathcal{L} = 2\mathcal{J}_+ \mathcal{J}_- + \mathcal{J}_- \mathcal{J}_-, \quad (26)$$

but new ordering gives (23):

$$\tilde{\mathcal{L}} = \mathcal{J}_+ \mathcal{J}_- + \mathcal{J}_- \mathcal{J}_-. \quad (27)$$

Here it is certainly possible to turn these expressions into each other by trivial rescaling of time variables. But this is already not the case when considering $\tilde{\mathcal{W}}^{(3)}$ -generators.

Indeed, the proper sum of currents in the $\tilde{\mathcal{W}}^3$ -case is:

$$(\mathcal{J}_+ \mathcal{J}_- \mathcal{J}_- + \mathcal{J}_- \mathcal{J}_+ \mathcal{J}_- + \mathcal{J}_- \mathcal{J}_- \mathcal{J}_+) + (\mathcal{J}_- \mathcal{J}_+ \mathcal{J}_+ + \mathcal{J}_+ \mathcal{J}_- \mathcal{J}_+ + \mathcal{J}_+ \mathcal{J}_+ \mathcal{J}_-) + \mathcal{J}_- \mathcal{J}_- \mathcal{J}_-. \quad (28)$$

Then terms in brackets are equal to each other under standard normal ordering and we obtain the formula (22).

In the case of $\tilde{\mathcal{W}}^{(3)}$ -generators in the second brackets the only term remains and the first two terms should be taken from the first brackets. But the term $\mathcal{J}_- \mathcal{J}_+ \mathcal{J}_-$ should be provided by additional restriction of the modes of bilinear combination of the right two currents $(\mathcal{J}_+ \mathcal{J}_-)_p$ to $p \geq -1$. Then this term corresponds to right ordering and leads to the second term in (20) and we obtain all items with unit coefficients as in (20). Thus we can rewrite the new ordering as follows:

$$\begin{aligned}
\tilde{\mathcal{W}}_p^{(K)} = \dagger(\partial\phi)^K \dagger_p = \dagger \mathcal{J}^K \dagger_p = & \sum_{\substack{p_1 + \dots + p_K = p \\ p_K \geq 0 \\ p_{K-1} + p_K \geq -1 \\ p_{K-2} + p_{K-1} + p_K \geq -2 \\ \dots}} \mathcal{J}_{p_1} \dots \mathcal{J}_{p_K}, \text{ with } p \geq 1 - K. \quad (29)
\end{aligned}$$

Its characteristic property is that $\dagger AB \dagger = \dagger A(\dagger B \dagger) \dagger$, but $\dagger AB \dagger \neq \dagger(\dagger A \dagger) B \dagger$.

The last comment concerns the possibility to reformulate the actual constraints (10) in terms of pure $\tilde{\mathcal{W}}^{(K)}$ -operators by appropriate shift of time variables. While this is trivial to do in the case of $K = 2$:

$$\tilde{\mathcal{W}}_n^{(2)}\{t_k\} - \frac{1}{2s_2} \frac{\partial}{\partial t_{n+2}} = \tilde{\mathcal{W}}_n^{(2)}\{t_k - \frac{1}{4s_k} \delta_{k,2}\} \quad (30)$$

things become worse already for $K = 3$. Namely, even to absorb derivative $\partial/\partial t_{p+3}$ into $\tilde{\mathcal{W}}_p^{(3)}$ one needs to perform a complicated recursive redefinition of times:

$$\tilde{t}_k = t_k + \Delta t_k, \quad (31)$$

$$\begin{aligned}
& 3s_3 \tilde{\mathcal{W}}_{p-2}^{(3)} \tilde{t} + 2s_2 \mathcal{W}_{p-1}^{(2)} \tilde{t} = \\
& = 3s_3 \tilde{\mathcal{W}}_{p-2}^{(3)}\{t\} + 2s_2 \mathcal{W}_{p-1}^{(2)}\{t\} + 3s_3 \sum_k k \Delta t_k \tilde{\mathcal{W}}_{p+k-2}^{(2)}\{t\} + \\
& + \left\{ 3s_3 \sum_{k+l=n+2} k(\Delta t_k) l(\Delta t_l) + 2s_2(n+1 + \Delta t_{n+1}) \right\} \frac{\partial}{\partial t_{p+n}}. \quad (32)
\end{aligned}$$

Thus it is clear that the most we can achieve is to adjust $\{\Delta t_k\}$ in such a way, that the expression in braces in the last term is equal to $-\delta_{n,1}$. Such adjusting is always possible, at least, in the iteration form

$$\begin{aligned}
\Delta t_1 &= 0, \\
\Delta t_2 &= -1/4s_2, \\
\Delta t_{n+1} &= -\frac{3s_3}{2(n+1)s_2} \sum_{k=2}^n k(n+2-k) \Delta t_k \Delta t_{n+2-k} \\
&\text{for } n > 1, \quad (33)
\end{aligned}$$

but it does not seem too illuminating.

4.3 Computation of $[\tilde{\mathcal{W}}_p^{(K)}, \tilde{\mathcal{W}}_q^{(K)}]$

To evaluate this commutator we shall make direct use of the definition (11) of $\tilde{\mathcal{W}}$ -operators and note that Λ and t_k are independent (and, therefore, commuting) variables. Thus

$$\begin{aligned}
\tilde{\mathcal{W}}_p^{(K+1)} \tilde{\mathcal{W}}_q^{(K+1)} e^{-trV} &= \tilde{\mathcal{W}}_p^{(K+1)} tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^K \Lambda^{q+K} \right\} e^{-trV} = \\
&= tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^K \Lambda^{q+K} \right\} \tilde{\mathcal{W}}_p^{(K+1)} e^{-trV} = \\
&= tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^K \Lambda^{q+K} \right\} tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^k \Lambda^{p+K} \right\} e^{-trV}, \tag{34}
\end{aligned}$$

so that

$$[\tilde{\mathcal{W}}_p^{(K+1)}, \tilde{\mathcal{W}}_q^{(K+1)}] e^{-trV} = \left[tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^K \Lambda^{q+K} \right\}, tr \left\{ \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^K \Lambda^{p+K} \right\} \right] e^{-trV} \tag{35}$$

(to simplify the formulae below we consider $\tilde{\mathcal{W}}_p^{(K+1)}$ instead of $\tilde{\mathcal{W}}_p^{(K)}$).

Our strategy in the derivation of (18) from the *r.h.s.* of (35) is as follows:

(i) Let us carry the second $(-\partial/\partial \Lambda_{tr})^K$ at the *r.h.s.* of (35) to the *left*. Neglecting all the commutators of $(-\partial/\partial \Lambda_{tr})^K$ with Λ^{q+K} , one obtains

$$\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)_{\alpha\beta}^K \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)_{\gamma\delta}^K (\Lambda^{q+K})_{\beta\alpha} (\Lambda^{p+K})_{\delta\gamma} e^{-trV}, \tag{36}$$

which cancels in the commutator (35). Therefore, only the terms $\left[(\Lambda^{q+K})_{\beta\alpha}, (-\partial/\partial \Lambda_{tr})_{\gamma\delta}^K \right]$ from the commutator are of interest, and we can consider

$$\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)_{\alpha\beta}^K \left\{ \left[(\Lambda^{q+K})_{\beta\alpha}, \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)_{\gamma\delta}^K \right] (\Lambda^{p+K})_{\delta\gamma} \right\} e^{-trV} - (p \leftrightarrow q) \tag{37}$$

instead of the *r.h.s.* of (35).

(ii) Expression in braces in (37) contains at most $K-1$ derivatives. Let us pull all these derivatives to the *right* and then use the relation

$$\left(-\frac{\partial}{\partial \Lambda_{tr}} \right)_{\mu\nu} e^{-trV} = (V')_{\mu\nu} e^{-trV} = \sum_k k t_k (\Lambda^{k-1})_{\mu\nu} e^{-trV}. \tag{38}$$

Then the factor $k t_k$ can be pulled to the very *left*, since t_k 's commute with all Λ 's and $\partial/\partial \Lambda_{tr}$'s.

(iii) What remains now from expressions in braces in (35) is a combination of powers of Λ 's which in fact has only two free indices $\beta\alpha$. Therefore, each item can be written as $(\Lambda^a)_{\beta\alpha}$ with some power a multiplied by some contribution of $tr \Lambda^b$ with various powers of b . All these traces can be shifted to the *right*. Then we use the fact that

$$(tr \Lambda^b) e^{-trV} = -\frac{\partial}{\partial t_b} e^{-trV}. \tag{39}$$

Again $\partial/\partial t_b$ can be carried to the left of all Λ 's and $\partial/\partial \Lambda'_{tr}$ s (but to the right of t'_k s which are arising at the step (ii)).

(iv) Thus we got rid of all contributions like $\partial/\partial\Lambda_{tr}$ and $tr\Lambda^b$ to the braces in (37). What remains is (a linear combination with “coefficients” made from kt_k and $\partial/\partial t_b$ of)

$$\left(-\frac{\partial}{\partial\Lambda_{tr}}\right)_{\alpha\beta}^K \Lambda_{\beta\alpha}^a e^{-trV} = \tilde{\mathcal{W}}_{a-K}^{(K+1)} e^{-trV}. \quad (40)$$

This is just the statement (18).

4.4 Two manifest examples

In order to illustrate this somewhat abstract consideration we present now two more explicit examples. The first one concerns Virasoro operators $\tilde{\mathcal{W}}^{(2)}$ — it is a quite trivial case. The expression from braces in (37) is now (note that $(\partial/\partial\Lambda_{tr})_{\alpha\beta} = (\partial/\partial\Lambda)_{\beta\alpha}$):

$$\left[(\Lambda^{q+1})_{\beta\alpha}, \left(-\frac{\partial}{\partial\Lambda_{tr}}\right)_{\gamma\delta}\right] (\Lambda^{p+1})_{\delta\gamma} = \sum_{a+b=q} (\Lambda^a)_{\beta\delta} (\Lambda^b)_{\gamma\alpha} (\Lambda^{p+1})_{\delta\gamma} = (q+1) (\Lambda^{p+q+1})_{\beta\alpha}, \quad (41)$$

and the *r.h.s.* of (35) becomes

$$[(q+1) - (p+1)] tr \frac{\partial}{\partial\Lambda_{tr}} \Lambda^{p+q+1} e^{-trV} = (p-q) \tilde{\mathcal{W}}_{p+q}^{(2)} e^{-trV}, \quad (42)$$

i.e.

$$[\tilde{\mathcal{W}}_p^{(2)}, \tilde{\mathcal{W}}_q^{(2)}] = (p-q) \tilde{\mathcal{W}}_{p+q}^{(2)}, \quad p, q \geq -1, \quad (43)$$

as it was expected. This example is not very representative as the steps (ii) and (iii) are in fact absent.

Another example of $\tilde{\mathcal{W}}^{(3)}$ -operators involves all the steps (i)-(iv). Expression in braces in (37) is now (note that $(\partial/\partial\Lambda_{tr})_{\alpha\beta}^2 = (\partial/\partial\Lambda)_{\epsilon\alpha} (\partial/\partial\Lambda)_{\beta\epsilon}$):

$$\begin{aligned} & - \left[(\Lambda^{q+2})_{\beta\alpha}, \left(\frac{\partial}{\partial\Lambda_{tr}}\right)_{\gamma\delta}^2 \right] (\Lambda^{p+2})_{\delta\gamma} = \\ & = - \sum_{a+b+c=q} [(\Lambda^a)_{\beta\epsilon} (\Lambda^b)_{\gamma\delta} (\Lambda^c)_{\epsilon\alpha} + (\Lambda^a)_{\beta\delta} (\Lambda^b)_{\epsilon\alpha} (\Lambda^c)_{\gamma\alpha}] (\Lambda^{p+2})_{\delta\gamma} + \\ & + \sum_{a+b=q+1} \left[(\Lambda^a)_{\beta\epsilon} (\Lambda^b)_{\gamma\alpha} \left(\frac{\partial}{\partial\Lambda}\right)_{\delta\epsilon} + (\Lambda^a)_{\beta\delta} (\Lambda^b)_{\epsilon\alpha} \left(\frac{\partial}{\partial\Lambda}\right)_{\epsilon\gamma} \right] (\Lambda^{p+2})_{\delta\gamma}. \end{aligned} \quad (44)$$

In order to complete the step (ii) we should carry the remaining derivatives $\partial/\partial\Lambda$ in the second sum to the right, so that the *r.h.s.* of (44) transforms into

$$\begin{aligned} & - \sum_{a+b+c=q} [(\Lambda^{a+c})_{\beta\alpha} tr \Lambda^{b+p+2} + (\Lambda^{a+c+p+2})_{\beta\alpha} tr \Lambda^b] - \\ & - \sum_{a+b=q+1} \sum_{c+d=p+1} [(\Lambda^{a+b+d})_{\beta\alpha} tr \Lambda^c + (\Lambda^{a+b+c})_{\beta\alpha} tr \Lambda^d] + \\ & + \sum_{a+b=q+1} \left[(\Lambda^a)_{\beta\epsilon} (\Lambda^{b+p+2})_{\delta\alpha} \left(\frac{\partial}{\partial\Lambda_{tr}}\right)_{\epsilon\delta} + (\Lambda^{a+p+2})_{\beta\gamma} (\Lambda^b)_{\epsilon\alpha} \left(\frac{\partial}{\partial\Lambda_{tr}}\right)_{\gamma\epsilon} \right]. \end{aligned} \quad (45)$$

Using the (38) we can turn it into

$$\begin{aligned}
& - \sum_{a+b=q} (a+1) [(\Lambda^a)_{\beta\alpha} tr \Lambda^{b+p+2} + (\Lambda^{a+p+2})_{\beta\alpha} tr \Lambda^b] - \\
& - 2(q+2) \sum_{a+b=p+1} (\Lambda^{a+q+1})_{\beta\alpha} tr \Lambda^b - 2(q+2) \sum_k kt_k (\Lambda^{p+q+k+2})_{\beta\alpha}.
\end{aligned} \tag{46}$$

Steps (iii) and (iv) are now trivial and we read off from (46):

$$\begin{aligned}
[\tilde{\mathcal{W}}_p^{(3)}, \tilde{\mathcal{W}}_q^{(3)}] &= 2(p-q) \sum_k kt_k \tilde{\mathcal{W}}_{p+q+k}^{(3)} - \\
& - \left\{ \sum_{a+b=q} (a+1) \left[\frac{\partial}{\partial t_{b+p+2}} \tilde{\mathcal{W}}_{a-2}^{(3)} + \frac{\partial}{\partial t_b} \tilde{\mathcal{W}}_{a+p}^{(3)} \right] + \right. \\
& + 2(q+2) \sum_{a+b=p+1} \frac{\partial}{\partial t_b} \tilde{\mathcal{W}}_{a+q-1}^{(3)} - (p \longleftrightarrow q) \Big\} = \\
& = 2(p-q) \sum_k kt_k \tilde{\mathcal{W}}_{p+q+k}^{(3)} + \\
& + \left[\sum_{a=0}^{p+1} (2p-q-2a) - \sum_{a=0}^{q+1} (2q-p-2a) \right] \frac{\partial}{\partial t_b} \tilde{\mathcal{W}}_{p+q-a}^{(3)}, p, q \geq -2.
\end{aligned} \tag{47}$$

For $p > q$ the item in square brackets on the *r.h.s.* can be equivalently rewritten as

$$3(p-1) \sum_{a+b=q+1} \frac{\partial}{\partial t_a} \tilde{\mathcal{W}}_{b+p-1}^{(3)} + \sum_{a=q+2}^{p+1} (2p-q-2a) \frac{\partial}{\partial t_a} \tilde{\mathcal{W}}_{p+q-a}^{(3)}. \tag{48}$$

For illustrative purposes let us show that operators $\tilde{\tilde{\mathcal{W}}}_p^{(3)} \equiv \tilde{\mathcal{W}}_p^{(3)} + \alpha(\partial/\partial t_{p+3})$ also form a closed set of constraints for $p \geq -2$. In accordance with (47), (48), it is enough to check that as soon as $p > q \geq -2$

$$\begin{aligned}
& \left[\frac{\partial}{\partial t_{p+3}}, \tilde{\mathcal{W}}_q^{(3)} \right] - (p \longleftrightarrow q) = 2(p-q) \sum_k kt_k \frac{\partial}{\partial t_{p+q+3}} + \\
& + 3(p-q) \sum_{a+b=q+1} \frac{\partial}{\partial t_b} \frac{\partial}{\partial t_{a+p+3}} + \sum_{b=0}^{p-q-1} (2p-3q-4-2b) \frac{\partial}{\partial t_{q+2+b}} \frac{\partial}{\partial t_{p+1-b}}.
\end{aligned} \tag{49}$$

The *l.h.s.* of this relation is easily evaluated with the help of explicit formula (20) for $\tilde{\mathcal{W}}_q^{(3)}$ and is equal to

$$\begin{aligned}
& 2(p-q) \sum_k kt_k \frac{\partial}{\partial t_{p+q+3}} + (p-q) \sum_{a+b=p+q+3} \frac{\partial^2}{\partial t_a \partial t_b} + \\
& + (p+3) \sum_{a+b=q+1} \frac{\partial^2}{\partial t_{a+p+2} \partial t_b} - (q+3) \sum_{a+b=p+1} \frac{\partial^2}{\partial t_{a+q+2} \partial t_b}.
\end{aligned} \tag{50}$$

It is a trivial arithmetic exercise to check that (50) coincides with the *r.h.s.* of (49) as soon as $p > q \geq -2$.

4.5 $\tilde{\mathcal{W}}$ -constraints in Kontsevich's like models

In order to add new colours to the picture of $\tilde{\mathcal{W}}$ -constraints let us present some details concerning their appearance in the framework of 1-matrix Kontsevich's models. Indeed, as shown in [6] the Ward identity (2) which has been applied above to the derivation of $\tilde{\mathcal{W}}^{(K)}$ -constraints in 2-matrix model may be represented in a form of $\tilde{\mathcal{W}}^{(K-1)}$ -constraint imposed on the partition function of 1-matrix model (1) itself. Following sect. 3.2 of ref.[6] we shall consider here only a simplified example, or what was called there a discrete counterpart of $\tilde{\mathcal{W}}$ -constraint in the model (1). Namely (see eq.(26) of ref.[6])

$$tr[M^q(\frac{\partial}{\partial M_{tr}})^K]\mathcal{F}\{t_k\} = \sum_{p \geq 1-K} tr M^{q-p-K} \tilde{\mathcal{W}}^{(K)} \mathcal{F}\{t_k\} . \quad (51)$$

for any function $\mathcal{F}\{t_k\}$ with the arguments

$$t_k = \frac{1}{k} tr M^{-k} , \quad (52)$$

$$t_0 = -tr \log M . \quad (53)$$

We shall demonstrate now that $\tilde{\mathcal{W}}^{(K)}$ -operators in the *r.h.s.* of (51) are exactly the same as $\tilde{\mathcal{W}}^{(K)}$'s of this paper defined through (11)

$$tr[(-\frac{\partial}{\partial \Lambda_{tr}})^{K-1} \Lambda^{p+K-1}] e^{-tr V\{\Lambda\}} = \tilde{\mathcal{W}}_p^{(K)} e^{-tr V\{\Lambda\}} . \quad (54)$$

In order to prove the equivalence of $\tilde{\mathcal{W}}$ -operators in (51) and (54) we take $\mathcal{F}\{t_k\} = e^{tr V}$, where

$$tr V = \sum_{k \geq 1} t_k tr \Lambda^k = -tr \log M - tr I + \sum_{k \geq 1} \frac{1}{k} tr M^{-k} tr \Lambda^k , \quad (55)$$

so that

$$e^{-tr V} = \det(M \otimes I - I \otimes \Lambda)^{-1} . \quad (56)$$

Then it remains to prove that (56) satisfies

$$tr[M^q(\frac{\partial}{\partial M_{tr}})^K] e^{-tr V} = \sum_{p \geq 1-K} tr M^{q-p-K} tr[(-\frac{\partial}{\partial \Lambda_{tr}})^{K-1} \Lambda^{p+K-1}] e^{-tr V\{\Lambda\}} \quad (57)$$

The *r.h.s.* of (57) may be further simplified: the sum over p is easily evaluated and gives

$$tr(M^q \otimes I)(I \otimes (-\frac{\partial}{\partial \Lambda_{tr}})^{K-1}) \frac{e^{-tr V}}{M \otimes I - I \otimes \Lambda} = tr[M^q(\frac{\partial}{\partial M_{tr}})] tr(\frac{\partial}{\partial M_{tr}})^{K-1} e^{-tr V} \quad (58)$$

so that (57) becomes a trivial identity

$$tr \left(M^q \frac{\partial}{\partial M_{tr}} \otimes I \right) \left(\left(\frac{\partial}{\partial M_{tr}} \right)^{K-1} \otimes I + I \otimes \left(-\frac{\partial}{\partial \Lambda_{tr}} \right)^{K-1} \right) \frac{1}{\det(M \otimes I - I \otimes \Lambda)} \equiv 0 \quad (59)$$

To make this proof a bit more transparent let us repeat it for the case of $N = 1$ (just numbers m and λ instead of matrices M and Λ):

$$m^q \left(\frac{\partial}{\partial m} \right)^K \frac{1}{m - \lambda} = \sum_{p \geq 1-K} m^{q-p-K} \left(-\frac{\partial}{\partial \lambda} \right)^{K-1} \lambda^{p+K-1} \frac{1}{m - \lambda} = m^q \left(-\frac{\partial}{\partial \lambda} \right)^{K-1} \frac{1}{(m - \lambda)^2} \quad (60)$$

Both sides of this relation are equal to $(K-1)! \frac{m^q}{(m-\lambda)^K}$ and, thus, coincide.

A really interesting question about Kontsevich's-like models (1) concerns the reformulation of actual Ward identities (2) in terms of appropriate time variables $T_k^{(K)}$, (in the case of $K = 2$ $T_k^{(2)}$ are defined by (3) instead of (52)-(53)). This is however beyond the scope of the present paper: the purpose of this section was to demonstrate that $\tilde{\mathcal{W}}$ -operators arise at least in two different contexts.

5 Conclusion

To conclude, we demonstrated that the Ward-identities of Kontsevich-like models, derived in [6], are enough to obtain a closed (and presumably) complete set of Ward identities (loop equations) in discrete 2-Hermitean matrix model. These loop-equations involve $\tilde{\mathcal{W}}$ -operators acting on one of potentials in the 2-matrix model, while the "highest" spin involved in these operators coincides with the power of another potential.³ This statement is the discrete counterpart of the suggestion of Gava and Narain [7], concerning the "asymmetric" (i.e. two potentials do not coincide) continuum limit of 2-matrix model, which is probably different from the symmetric limit, originally examined in [2].

We demonstrated also that these unconventional $\tilde{\mathcal{W}}$ -operators arise at least at two slightly different places. First, if we study an analogue of Kontsevich's model with potential $W[X]$ of M -th power (or rather what was called in [6] its discrete "analogue"), then operators $\tilde{\mathcal{W}}^{(M-1)}$ arise. Second, if just the same X -integral is treated as the constituent of 2-matrix model, then operators $\tilde{\mathcal{W}}^{(M)}$ arise. Note that in these two examples the "spins" of $\tilde{\mathcal{W}}$ -operators are different. Of course, their meanings are different too, as well as the time-variables, through which they are expressed: these are really two *different* examples where *the same* $\tilde{\mathcal{W}}$ -operators seem to arise.

Detailed study of $\tilde{\mathcal{W}}$ -algebras, their relations to conventional $\mathcal{W}$ -algebras, their continuum limits (in the matrix-model sense) etc. deserve further analysis, which is beyond the scope of this Letter.

Acknowledgements

We benefited a lot from illuminating discussions on the subject with E.Corrigan, E.Gava, K.Narain, A.Niemi.

³So one can look at these constraints as at Ward identities of proper 1-matrix model (in terms of variable Λ). It differs from "true" 2-matrix model $\mathcal{W}$ -constraints depending on two sets of times, see the example of this in [11].

We are grateful to the hospitality of the Research Institute for Theoretical Physics (TFT) at Helsinki University where some pieces of this work have been done. A.Mor. is also very much indebted for the hospitality of Dept. of Math. Sci. of Durham University. The work of A.Mar. and A.Mir. was partially supported by the Research Institute for Theoretical Physics (TFT) at Helsinki University while that of A.Mor. by the UK Science and Engineering Council through the Visiting Fellowship program and by NORDITA.

References

1. V.Kazakov Mod.Phys.Lett. A4 (1989) 2125
E.Brezin, V.Kazakov Phys.Lett. B236 (1990) 144
M.Douglas, S.Shenker Nucl.Phys. B335 (1990) 635
D.Gross, A.Migdal Phys.Rev.Lett. 64 (1990) 127
2. M.Fukuma, H.Kawai, R.Nakayama Int.J.Mod.Phys. A6 (1991) 1385
R.Dijkgraaf, E.Verlinde, H.Verlinde Nucl.Phys. B356 (1991) 574
3. A.Gerasimov et.al. Nucl.Phys. B357 (1991) 565
4. Yu.Makeenko et.al. Nucl.Phys. B356(1991) 574
5. M.Kontsevich Funk.Anal. i Priloz. 25 (1991) 50
6. A.Marshakov et al. "On equivalence of topological and quantum $2d$ gravity", Phys.Lett. B, in press
7. E.Gava, K.Narain Phys.Lett. B263 (1991) 213
8. M.L.Mehta Comm.Math.Phys. 79 (1981) 327
9. A.Mironov, A.Morozov Phys.Lett. B252 (1990) 47
10. J.Ambjorn, J.Jurkiewicz, Yu.Makeenko Phys.Lett. B251 (1990) 517
11. A.Marshakov, A.Mironov, A.Morozov Phys.Lett. B265 (1991) 99